



Environmental Microbiology

Control potential of *Meloidogyne javanica* and *Ditylenchus* spp. using fluorescent *Pseudomonas* and *Bacillus* spp.



Marcielly F. Turatto^a, Fernanda dos S. Dourado^b, Jerri E. Zilli^b, Glória R. Botelho^{c,*}

^a UFSC Centro Curitibanos, Curitibanos, SC, Brazil

^b Embrapa – Agrobiologia, Seropédica, RJ, Brazil

^c UFSC Centro Curitibanos, Curitibanos, SC, Brazil

ARTICLE INFO

Article history:

Received 1 July 2016

Accepted 17 March 2017

Available online 30 August 2017

Associate Editor: Fernando Andreote

Keywords:

Allium sativum L.

Glycine max (L.) Merrill

Biocontrol

Plant parasitic nematodes

Rhizobacteria

ABSTRACT

Plant Growth Promoting Rhizobacteria (PGPR) have different mechanisms of action in the development of plants, such as growth promotion, production of phytohormones and antibiotic substances and changes in root exudates. These help to control plant diseases. In order to evaluate the potential of microorganisms in the control of *Meloidogyne javanica* and *Ditylenchus* spp., five rhizobacteria isolated from rhizosphere of garlic cultivated in the Curitibanos (SC) region were tested. Hatching chambers were set on Petri dishes, in which were added 10 mL of bacterial suspension and 1 mL of *M. javanica* eggs suspension, at the rate of 4500, on the filter paper of each chamber. The same procedure was performed with 300 juvenile *Ditylenchus* spp. The experimental design was completely randomized, with four replications. The evaluations were performed every 72 h for nine days. The antagonized population of nematodes was determined in Peters counting chamber, determining the percentage hatching (for *M. javanica*) and motility (for *Ditylenchus* spp). Isolates CBSAL02 and CBSAL05 significantly reduced the hatching of *M. javanica* eggs (74% and 54.77%, respectively) and the motility of *Ditylenchus* spp. (55.19% and 53.53%, respectively) *in vitro*. Isolates were identified as belonging to the genera *Pseudomonas* (CBSAL05) and *Bacillus* (CBSAL02).

© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Some of the plant crops commodity cultivated in Curitibanos region (SC) are garlic (*Allium sativum*) and soybean (*Glycine max* (L.) Merrill). Those two crops are used in the main rotation system adopted by local farmers. In the case of garlic production,

Curitibanos ranks as the leading producer in the state of Santa Catarina. Soybean production in the region is low; however, the genetic potential of the crop generates a secure market and serves as an alternative crop to increase the income of the producer.

As with all crops, garlic and soybeans are also susceptible to a variety of pests and diseases. Among biotic factors,

* Corresponding author at: UFSC Centro Curitibanos, Rod. Ulysses Gaboardi s/n, km.03, CEP 89520-000, Curitibanos, SC, Brazil.

E-mail: gloria.botelho@ufsc.br (G.R. Botelho).

<https://doi.org/10.1016/j.bjm.2017.03.015>

1517-8382/© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

nematodes cause significant drop in production. These organisms parasitize both the roots and the aerial parts of the plants.

The main nematode parasite in garlic is *Ditylenchus dipsaci*, which is distributed in all production regions of Brazil.¹ One of the factors hindering its control is the anhydrobiosis ability of the fourth instar juvenile.² In addition, few chemicals are released by the Ministry of Agriculture Livestock and Food Supply to control this agricultural pest.³

On the other hand, the nematodes of the genus *Meloidogyne* cause the most damage to soybean.⁴ Due to monoculture or rotation with plant species that are also the host, coupled with low ground cover, the genus finds a favorable environment for its development.

Currently, control of nematodes is accomplished through the use of nematicides that are high-priced products and have residual action, as well as high toxicity to the environment and soil microorganisms.⁵ Because of the possible negative impacts that it can cause⁶ and restrictions in the use of these products, biological control can become a great ally in the control of these plant parasites.⁷

An alternative to chemical control is the use of antagonist microorganisms. The Plant Growth Promoting Rhizobacteria (PGPR) can be powerful agents for biological control of plant parasitic nematodes. *Bacillus* and fluorescent *Pseudomonas* group are among the most studied and with the highest correlation to soil suppressiveness.⁸ Rhizobacteria have several mechanisms of action for the plant disease control.⁹ The main mode of action of this group of bacteria is through the production of enzymes, antibiotics, siderophores, changes in root exudates, and induced resistance, among others.¹⁰ According to Becker et al.,¹¹ changes in root exudates can inhibit the hatching of eggs in nematodes, or even reduce the attractiveness of these to the roots of plants. Freitas et al.¹² confirmed the potential of *Pseudomonas fluorescens* isolates to control nematodes by achieving a 75% control of *Heterodera schachtii*. The same isolates also controlled *Meloidogyne* spp. and *Radopholus similis* in maize, banana and tomato. Isolates of the same bacterial species have been reported as effective in the inhibition of *M. incognita* on tobacco, when inoculated into the soil with *Trichoderma harzianum*, and the action resembles the nematicides Furadan and phorate.¹³

In this paper, the objective was to evaluate the effect of rhizobacteria, belonging to Rhizobacteria collection of Laboratório de Microbiologia da UFSC Centro de Curitiba, in the control of *Meloidogyne javanica* and *Ditylenchus* spp.

Materials and methods

Effect of rhizobacteria isolates in the hatching of eggs of *M. javanica* in vitro

Five isolates of rhizobacteria (CBSAL02, CBSAL05, CBSAL14, CBSAL18 and CBSAL21) from Rhizobacteria Collection of the Laboratório de Microbiologia from Universidade Federal de Santa Catarina Centro Curitiba were selected. The rhizobacteria were grown in Petri dishes containing solid King B medium and subsequently incubated for 24 h at 28 °C. After

growth, the suspension of these isolates was held in sterile distilled water with the aid of Drigalski strap, under aseptic conditions. Each suspension was placed into sterile glass bottles. The optical density of the bacterial suspensions was adjusted to 0.2 optical density (wavelength 625 nm).

The eggs of *M. javanica* used in the experiment were provided by Dr. Bruno Barbosa Unesp Jaboticabal (SP). Hatching chambers were assembled according to the methodology proposed by Alves et al.,¹⁴ in which were added 10 mL of each bacterial suspension. Then, onto the filter paper was added 1 mL of the egg suspension of *M. javanica*, containing approximately 4500 eggs. As a control treatment, only sterile water with *M. javanica* eggs was used. The chambers were conditioned at a temperature of 29 °C for a period of nine days. The statistical design was completely randomized, with six treatments and four replications, each of which has a hatching chamber. For every 72 h (three days), aliquots were removed from the suspensions of the chambers for evaluation of the degree of inhibition of egg hatching by each bacterial suspension.

The population of nematodes hatched, which was retained on the filter paper, was determined with the aid of an optical microscope and Peters counting chamber. Nematodes recovered below the paper were recorded as not antagonized. For counting was determined the hatched individuals in the control (100% hatching) and then, in the other treatments. The percentage of hatching was calculated by comparison with the number in the control. The volume of bacterial suspension removed was replaced individually in each chamber, after the withdrawal according to each treatment.

Effects of rhizobacteria isolates on the motility of *Ditylenchus* spp.

The preparation of the inoculum followed the same methodology described above.

The population of nematodes was extracted from garlic plants contaminated by the nematode samples. The methodology used was flotation–centrifugation in sucrose solution with kaolin.¹⁵ After extraction, individual nematodes were placed in Petri dishes containing 20% formaldehyde. Later, the nematodes were identified with the help of the identification key proposed by Mekete et al.¹⁶ After being identified as belonging to *Ditylenchus* genus, the population was established as 300 nematodes/mL in the extracted suspension, regardless of the stage of development of the individual.

Hatching chambers were prepared and to it 10 mL of each bacterial suspension was added. Then, to the filter paper, 1 mL of *Ditylenchus* spp. suspension containing approximately 250 individuals, at different stages of development, was added. As a control treatment, only sterile water added to the filter paper with nematodes was used. The chambers were placed in 29 °C, for a period of nine days. The statistical design was completely randomized, with six treatments and four replications, with each plot having a hatching chamber.

At intervals of 72 h, samples were taken from the suspensions in the chambers to evaluate the degree of antagonism of each isolate.

The population of antagonized nematodes retained on the filter paper was determined with the aid of an optical

Table 1 – Inhibition of egg hatching in *Meloidogyne javanica* by the PGPR isolates.

	Control	CBSAL21	CBSAL05	CBSAL18	CBSAL02	CBSAL14
Rep1	3060	2160	1370	2560	760	1490
Rep2	2630	2080	1250	2480	680	1370
Rep3	2740	2160	1220	2250	760	1450
Rep4	2780	2010	1230	2240	720	1370
Average	2802.5 ^a	2102.5 ^c	1267.5 ^d	2382.5 ^b	730 ^e	1420 ^d
% of hatching	100%	75%	45.23%	114%	26%	50.67%
% of inhibition	0%	25%	54.77%	-14%	74%	49.33%

Means followed by the same letter do not differ. Tukey test at 5% probability.

Table 2 – Inhibition of the motility of different juvenile stages of *Ditylenchus* spp. by the PGPR isolates.

	CBSAL18	CBSAL14	CBSAL21	CBSAL02	CBSAL05	Control
Rep1	24	54	29	33	30	57
Rep2	53	32	31	21	29	64
Rep3	37	44	27	27	23	57
Rep4	40	53	23	27	30	63
Total	154 ^b	183 ^{ab}	110 ^b	108 ^b	112 ^b	241 ^a
Average	5.92186 ^b	6.72967 ^{ab}	5.23623 ^b	5.17986 ^b	5.28386 ^b	7.75923 ^a
% of inhibition	36.09%	24.09%	54.36%	55.19%	53.53%	0.00%

Means followed by the same letter do not differ. Tukey test at 5% probability.

microscope and Peters counting chamber. Nematodes recovered below the paper were recorded as not antagonized. To calculate the degree of antagonism, the number of individuals in the control (considered 100% not antagonized) and then the number in the other treatments were determined. The percentage of antagonism was calculated by comparing with the number in the control. The volume of bacterial suspension removed was replaced individually in each chamber after the withdrawal according to each treatment.

Statistical analysis

The data relating to *M. javanica* hatching test and motility of *Ditylenchus* spp. were subjected to analysis of variance and the means were compared by Tukey test at 1% and 5% probability to the statistical program Assistat version 7.7.

Amplification and sequencing of 16S rRNA gene

The bacteria were initially grown in King B medium for 24 h and then subjected to DNA extraction using the Wizard Clean up extraction kit following the manufacturer's recommendations. Amplification was performed in a final volume of 50 μ L, containing approximately 50 ng of template DNA, 1 \times buffer, 1.75 mM Mg Cl₂, 0.25 mM dNTP each, primers 27F and 1492R and 0.20 mM each Taq polymerase 1.5 U. Sequencing was performed using sense and antisense primers in equipment of Applied Biosystems 3500. The assembly of the contigs was done using BioNumerics 7.0 program.

The sequences obtained were subjected to a similarity analysis in the NCBI database (National Center for Biotechnology Information) using the BLAST tool (Basic Local Alignment Search Tool).¹⁷

Results

Effect of isolates of rhizobacteria in the hatching of eggs of *M. javanica* and the motility of *Ditylenchus* spp. in vitro

To calculate the percentage of control, comparison index was used. The control treatment that received no bacteria was considered as 100% hatching of *M. javanica* eggs or 0% of control. The isolates CBSAL02, CBSAL05, CBSAL14 and CBSAL21 presented at the end of the evaluation averages significantly higher than other treatments, taking into account the overall average of the treatments. These showed 74% control rates, 54.77%, 49.33% and 25%, respectively (Table 1). Isolated CBSAL18 showed an inhibition percent significantly lower than the control, suggesting no antagonistic effect.

Regarding the results of the tests inhibiting the motility of *Ditylenchus* spp., isolates CBSAL02, CBSAL05, CBSAL18, and CBSAL21 statistically differed from the control, while the isolate CBSAL14 statistically matched other treatments and the control (Table 2). The isolates CBSAL02, CBSAL21, CBSAL05 and CBSAL18 exhibited control of 55.19%, 54.36%, 53.53% and 36.09%, respectively. CBSAL14 presented a percentage of 24.09% (Table 2); however, compared to the control, the difference was not considered significant.

Identification of rhizobacteria isolates

Comparison of 16S rRNA sequences of five isolates (CBSAL02, CBSAL05, CBSAL14, CBSAL18 and CBSAL21) showed that three isolates (CBSAL05, CBSAL18 and CBSAL21) belonged to the genus *Pseudomonas*, with 99% similarity (Table 3). Other

Table 3 – Identification of the PGPR isolates by 16S rRNA gene sequencing.

Isolates	Identification	ID/NCBI	Similarity (%)	E-value
CBSAL02	<i>Bacillus</i> sp.	AB733561.1	99	0.0
CBSAL05	<i>Pseudomonas</i> sp.	KJ601736.1	99	0.0
CBSAL14	<i>Klebsiella</i> sp.	KT860061.1	99	0.0
CBSAL18	<i>Pseudomonas</i> sp.	KT825738.1	99	0.0

isolates were identified as *Bacillus* (CBSAL02) and *Klebsiella* (CBSAL14); both also showed 99% similarity (Table 3).

Discussion

Four of the isolates tested belonged to *Pseudomonas* and *Bacillus* genera, which are commonly described as inhibitors of various pathogens.^{9,18,19} The *Klebsiella* genus is described endophytic, diazotrophic and a plant growth promoter as well.²⁰ Information regarding its effect on plant pathogens was found in the literature.

Among the three isolates identified as *Pseudomonas*, one inhibited over 50% the two nematodes, while the other two showed inhibition percentage at around 50% to only one of the parasitic nematodes. The results indicated an antagonistic action of the isolates to the nematodes, suggesting potential control. It was noted that two of the five isolates (CBSAL02 and CBSAL05) had percentages of control higher than 50% to the two nematodes examined. The control ability of CBSAL21 to *Ditylenchus* spp. was more than double of that observed for *M. javanica* (54, 36% and 25% respectively). A reverse condition was observed for CBSAL14, wherein the percentage for *M. javanica* was 49.33% and 24.09% for *Ditylenchus* spp. For CBSAL18, there was inhibition of 36% of *Ditylenchus* spp., while for *M. javanica*, there was no effect on hatching. These results suggest specific action of some rhizobacteria to taxonomic groups of nematodes.

The results are consistent with many previous studies to assess the effect of rhizobacteria on the control nematodes. Alves et al.¹⁴ demonstrated through *in vitro* experiments the potential of different isolates of rhizobacteria, with regard to the motility and ovicidal action of *M. javanica*, *M. incognita* and *P. zeae*. Naves et al.²¹ observed that endophytic bacteria filtered suspensions were able to significantly reduce the mobility, and consequently increase mortality, as well as the hatching of the young second stage of *M. javanica*.

Some authors have reported rhizobacterial antagonistic action to be dependent on the production ability of secondary metabolites, which could explain the effect variance observed in some isolates by the production or not of specific compounds. Studies by Stirling²² confirm that some rhizobacterial isolates are able to synthesize toxic metabolites, affecting the movement and the emergence of the young in different kinds of nematodes.

Extracellular enzymes are the most studied rhizobacterial metabolites with inhibitory action against pathogens. This action has also been studied in other microorganisms. Khan et al.²³ reported the production of proteases and chitinases by an isolated fungus *Paecilomyces lilacinus*. Those substances were observed to operate together, destroying the lipid layer that is essential for the development and maintenance of

nematode in the egg; in addition, they caused the hydrolysis of egg chitin layer and affected the integrity of the vitelline layer, thus having an impact on both development and hatching.

It is suggested that the inhibitory action of rhizobacterial isolates may be linked to the production of enzymes, such as chitinase and other cell wall degrading substances. These would assist in the degradation of the wall of *M. javanica* eggs and, subsequently, impair the development of that nematode, as well as *Ditylenchus* spp. Some PGPR produce these lytic enzymes that break down the walls of the eggs of species of *Meloidogyne*, delay the onset of second stage juveniles (J2), cause the death of adult nematodes or interfere with their host plant recognition process.²⁴

The recognition of the plant host is the primary factor in the infection by nematode. The plant identification is performed through the root exudates, where nematodes will migrate to the infection site. Rhizobacteria as those of the genus *Pseudomonas* possess the ability to alter root exudates, thus hampering the recognition of the feeding site by the nematode present in the soil.²⁵ Siddiqui and Mahmood²⁶ and Tian et al.²⁷ observed that the interference in the recognition process by exudate changes was caused by endophytic bacteria as a control mechanism.

The action of rhizobacterial isolates in plants has been reported in several studies. Araújo et al.²⁸ demonstrated the efficacy of *Bacillus* spp. and *Pseudomonas* spp. in the control of *R. similis*, which causes root necrosis in banana plants. They also pointed out the prevalence of these bacterial genera in the rhizosphere of different plants. De Souza Júnior et al.²⁹ evaluated the effect caused by the mixing of rhizobacterial isolates for the control of *Meloidogyne graminicola* in rice seedlings grown from microbiolized seeds. In this study, the authors observed a reduction in the number of galls, number of eggs and reproduction of *M. graminicola*.

Five bacterial isolates were inoculated in lettuce pre-inoculated seedlings grown in substrate (vermiculite:sand – 2:1) with *M. javanica*. However, due to unfavorable environmental factors in the development of nematode, there was no gall formation.

Tested rhizobacteria also exhibited plant growth promoting activity *in vitro* and *in vivo*, indicating its potential use in production and plant protection.³⁰

The bacterial isolates CBSAL02, CBSAL05, CBSAL14 and CBSAL21 provided ovicidal action *M. javanica*. Aside from CBSAL14, all the isolates showed potential control to *Ditylenchus* spp. when compared to the control treatment. The results indicate that two isolates (CBSAL02 and CBSAL05) exhibited strong potential for use as biological control agents for both the nematodes. Detailed studies of the action of these isolates on nematodes should be performed, as well as evaluations on their potential for plant protection.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors thank Professor Jaime Maia dos Santos and Dr. Bruno F. F. Barbosa, Unesp Jaboticabal, for the assignment regarding the nematodes and technical support.

REFERENCES

- Charchar JM, Tenente RCV, Aragão FAS. Resistência de cultivares de alho a *Ditylenchus dipsaci*. *Nematol Bras*. 2003;27:179–184.
- Pinheiro JB, De Carvalho ADF, Pereira RB, Rodrigues C da S. *Nematoides na cultura do alho e cebola*. Brasília-DF, Góias: Embrapa Hortaliça; 2014.
- MAPA. *Agrofit*; 2014. Available at: http://agrofit.agricultura.gov.br/agrofit_cons/principal_agrofit_cons Accessed 28.09.14.
- Kimati H, Amorim L, Rezende JAM, Bergamin Filho A, Camargo LEA, eds. *Manual de Fitopatologia. Doenças das Plantas Cultivadas*. vol. 2, 4th ed. São Paulo: Ceres; 2005.
- Trigiano RN, Windham MT, Windham AS. *Fitopatologia: conceitos e exercícios de laboratório*. In: Noe JP, ed. *Nematoides parasitas de plantas*. 2nd ed. São Paulo: Artmed; 2010:83–96.
- Michereff SJ. *Fundamentos de Fitopatologia*. Recife: UFRPE; 2001.
- Dos Santos JM, Soares PLM, Barbosa BFF. *Curso de Atualização em Nematologia*. CD-ROM. Curitiba, SC: UFSC; 2013.
- Bettiol W, Ghini R, Mariano RRL, et al. Supressividade a fitopatógenos habitantes do solo. In: Bettiol W, Morandi MAB, eds. *Biocontrole de doenças de plantas: usos e perspectivas*. Jaguariúna: Embrapa Meio ambiente; 2009: 187–190.
- Botelho GR, Mendonça-Hagler LC. Fluorescent *Pseudomonas* associated with the rhizosphere of crops – an overview. *Braz J Microbiol*. 2006;37:401–416.
- Ludwig J, Moura AB, Gomes CB. Potencial da microbiolização de sementes de arroz com rizobactérias para o biocontrole do nematoide das galhas. *Trop Plant Pathol*. 2013;38(3): 264–268.
- Becker JO, Zavaleta-Mejia E, Colbert SF, et al. Effect of rhizobacteria on root-knot nematodes and gall formation. *Phytopathology*. 1988;78:1466–1469.
- Freitas LG, Neves WS, Fabry CSF, et al. Isolamento e seleção de rizobactérias para controle de nematoides formadores de galha (*Meloidogyne* spp.) na cultura do tomateiro. *Nematol Bras*. 2005;29(2):215–220.
- Khan MR, Haque Z. Soil application of *Pseudomonas fluorescens* and *Trichoderma harzianum* reduces root-knot nematode, *Meloidogyne incognita*, on tobacco. *Phytopathol Mediterr*. 2011;50:257–266.
- Alves GCS, dos Santos JM, Soares PPLM, de Jesus FG, de Almeida EJ, Thuler RT. Avaliação *in vitro* do efeito de rizobactérias sobre *Meloidogyne incognita*, *M. javanica* e *Pratylenchus zea*. *Arq Inst Biol*. 2011;78(4): 557–564.
- Coolen WA, D’Herde CJA. *Method for the Quantitative Extraction of Nematodes from Plant Tissue*. Belgium, Gent: State Agricultural Research Center, Gent; 1972.
- Mekete T, Dababat A, Sekora N, Akyazi F, Abebe E, comps. Identification key for agriculturally important plant-parasitic nematodes. Prepared for the International Nematode Diagnosis and Identification Course 2012 – A manual for nematology. Mexico: CIMMYT; 1972.
- Altshul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215:403–410.
- Lanna Filho R, Ferro HM, De Pinho RSC. Controle biológico mediado por *Bacillus subtilis*. *Rev Tróp Ciênc Agrár Biol*. 2010;4(2):12–20.
- Sivasakthi S, Usharani G, Saranraj P. Biological potentiality of plant growth promoting bacteria (PGPR) – *Pseudomonas fluorescens* and *Bacillus subtilis*: a review. *Afr J Agric Res*. 2014;9(16):1265–1277.
- Moreira FM de S. Bactérias diazotróficas associativas: diversidade, ecologia e potencial de aplicações. *Comun Sci*. 2010;1(2):74–99.
- Naves RL, Campos VP, Souza RM. Filtrados de culturas bacterianas endofíticas na motilidade, mortalidade e eclosão de juvenis de segundo estágio de *Meloidogyne javanica*. *Fitopatol Bras*. 2004;29(4):384–388.
- Stirling GR. *Biological Control of Plant Parasitic Nematodes: Progress, Problems and Prospects*. Wallingford: CAB International; 1991.
- Khan A, Williams KL, Nevalainen HKM. Effects of *Paecilomyces lilacinus* protease and chitinase on the eggshell structures and hatching of *Meloidogyne javanica* juveniles. *Biol Control*. 2004;31:346–352.
- Spiegel Y, Mor M, Sharon E. Attachment of *Pasteuria penetrans* spores to the surface of *Meloidogyne javanica* second-stage juveniles. *J Nematol*. 1996;28:328–333.
- De Melo IS, De Azevedo JL, eds. *Ecologia microbiana*. Jaguariúna: Embrapa Meio Ambiente; 1998.
- Siddiqui ZA, Mahmood I. Role of bacteria in the management of plant parasitic nematodes: a review. *Bioresour Technol*. 1999;69:167–179.
- Tian B, Yang J, Zhang KQ. Bacteria used in the biological control of plant-parasitic nematodes: populations, mechanisms of action, and future prospects. *FEMS Microbiol Ecol*. 2007;61:197–213.
- Araújo KS, Peixoto CC, Da Silva AC, Cardoso KGV, Da Silva HSA, Trindade AV. Avaliação e identificação de *Pseudomonas* sp. e *Bacillus* sp. dois isolados de rizobactérias antagonicas a *Radopholus similis*. In: *Jornada Científica – Embrapa Mandioca e Fruticultura*.; 2010. Available at: <http://ainfo.cnptia.embrapa.br/digital/bitstream/item/26342/1/071-Kalianel-Aldo-ok.pdf> Accessed 05.04.16.
- De Souza Júnior IT, Moura AB, Schafer JT, Corrêa BO, Gomes CB. Biocontrole da queima-das-bainhas e do nematoide-das-galhas e promoção de crescimento de plantas de arroz por rizobactérias. *Pesq Agropecu Bras*. 2010;45(11):1259–1267.
- Leoncio MR, Botelho GR, Orsi B, Soares CRFS, De Armas RD, Lovato PE. Isolamento e caracterização de rizobactérias indutoras de crescimento vegetal no alho (*Allium sativum*). In: *X Reunião Sul-Brasileira de Ciência do Solo: Fatos e Mitos em Ciência do Solo*. 2014. Available at: <http://www.sbcs-nrs.org.br/xrsbcs/docs/trab-3-4758-72.pdf> Accessed 05.04.16.