



## **Effectiveness of *Azospirillum brasilense* Inoculants to Wheat (*Triticum aestivum*) in the Micro-region of Curitibanos (SC)**

**William Gilberto Balbinot<sup>1</sup>, André Luis Gordechuk<sup>1</sup>,  
Géssica Rogaleski Eutrópio<sup>1</sup>, Cibele Medeiros<sup>2</sup> and Glória Regina Botelho<sup>1\*</sup>**

<sup>1</sup>Universidade Federal de Santa Catarina (UFSC), Centro de Ciências Rurais, Campus de Curitibanos, Rodovia Ulisses Gaboardi, Km.03 – 89520-000, Curitibanos, SC, Brazil.

<sup>2</sup>Grupo Vítia, Biosoja Agrociência, Av. Marginal Esquerda, 2000, Via Anhanguera, Km 383 - 14.600-000, São Joaquim da Barra, SP, Brazil.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors WGB, ALG and GRE performed the experimental and statistical analysis. Authors CM and GRB wrote the manuscript. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JEAI/2020/v42i130450

#### Editor(s):

- (1) Dr. Suleyman Korkut, Professor, Division of Wood Mechanic and Technology, Department of Forest Industrial Engineering, Duzce University, Turkey.  
(2) Dr. Vincenzo Tufarelli, Professor, Department of DETO - Section of Veterinary Science and Animal Production, University of Bari "Aldo Moro", Italy.

#### Reviewers:

- (1) Viviane Cristina Modesto, Sao Paulo State University, Brazil.  
(2) Clovis Arruda De Souza, Santa Catarina State University, Brazil.  
(3) Dahiru Ibrahim, National Research Institute for Chemical Technology, Nigeria.  
(4) Moataz Eliw Mostafa, Al-Azhar University, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/53471>

**Original Research Article**

**Received 28 November 2019**

**Accepted 03 January 2020**

**Published 15 February 2020**

### **ABSTRACT**

**Aims:** This work aimed to evaluate the effectiveness of *A. brasilense* inoculants to the development of wheat at the southern part of Brazil.

**Study Design:** The experimental design was randomized block with twelve treatments, containing two liquid inoculants with the *A. brasilense* strains Ab-V5 and Ab-V6, with or without nitrogen fertilization and five replicates. The plot area was 4m per 6m and thirty-two rows. The seeds inoculation was performed according to the manufacturer's recommendation, and manual sowing.

\*Corresponding author: E-mail: [gloria.botelho@ufsc.br](mailto:gloria.botelho@ufsc.br), [grbotelho@hotmail.com](mailto:grbotelho@hotmail.com);

The top-dressing N fertilization was Urea (Super N- 45%N) at 20 days after emergence (DAE), at 120 kg ha<sup>-1</sup> (full dosage) or 60 kg ha<sup>-1</sup> (half dosage).

**Place and Duration of Study:** The experiment was performed in a farm in Curitiba county in Santa Catarina state, Brazil. The experiment was carried out on July to November 2016.

**Methodology:** At 45 DAE, it was performed flag leaves N content Tedesco, et al. [1]. At 115 DAE, it was performed dry shoot weight, plant height, ear sizes, grain N contents and grain yield. The results were submitted to variance analysis (ANOVA) and media compared by Scott-Knot's test at 5% of significance.

**Results:** There was no statistical difference for shoot dry weight, plant height, ear size and yield. The N leaf content was greater with Ab-V5 inoculation and half N dosage (HC5 - 109% higher than the control). The N grain content was greater with the two strains (inoculant B) without, half and full N dosage (WC56 - 51%, HCB56 - 76% and CB56 - 65%, respectively).

**Conclusion:** *A. brasilense* strains had the ability to increase wheat N accumulation with lower N fertilizing, suggesting their potential as growth inducers, emphasizing the importance of further studies to confirm and understand the mechanisms involved.

**Keywords:** *Inoculation; N content; plant growth promotion; rhizobacteria; Poaceae.*

## 1. INTRODUCTION

In 2019, wheat production reached 5.20 million tons in Brazil [2], about 7% lower than in the previous harvests. This value corresponds to approximately 50% of the cereal consumption, around 10 tons [3]. The estimate for the crop projected reduction of area and production, due to environmental factors, costing, price and production of the last harvest [3,4]. This trend was confirmed, especially in the southern states of the country (Rio Grande do Sul, Paraná and Santa Catarina states) that are the main producers. The decline in area and production of wheat in these states was approximately 10% and 26%, respectively [4]. Santa Catarina recorded a fall of 14% [4] in its agricultural regions, including Curitiba county, one of the four most important. It is clear that the adoption of strategies to encourage cereal production is necessary, especially those that could reduce production costs and ensure the sustainability of agricultural systems.

Nitrogen plays an important key role in wheat yield. The availability of this nutrient directly influences crop yield [5]. Depending on the content of SOM (Soil Organic Matter) and the previous crop, 60-80 kg N ha<sup>-1</sup> are recommended [5]. In several crops, including wheat, urea is the most commonly used nitrogen fertilizer and represents about 40% of the production cost. Besides burdening the budget, it can lead to environmental disturbances [6]. The biological N fixation (BNF) has been an important and viable alternative for several crops [7,8,9,10]. The technology of N-fixing bacteria inoculation is quite widespread [9,11]. Several bacteria, such as *Azospirillum brasilense*, have been studied

because it induces plant growth by mechanisms as BNF, phytohormone production, among others [7,12,13]. Strains Ab-V5 and Ab-V6 from *A. brasilense* increased corn and wheat yields by 27% and 31%, respectively [14]. Currently, the strains are part of several commercial inoculants and it is necessary to comprehend their efficiency in different edaphoclimatic conditions. In Curitiba county, located in Santa Catarina state, inoculation is used for crops, such as soybean and maize. However, there is no information about its application to wheat. Based on it, the objective was to evaluate the effectiveness of *A. brasilense* on the development of wheat at this region.

## 2. MATERIALS AND METHODS

### 2.1 Description of the Experimental Area

The experiment was sown at a farm in Curitiba County, Santa Catarina, Brazil. This is located at latitude 27°16'58" south, longitude 50°35'04" west and altitude of 987 m. According to the Köppen classification, the climate is Cfb featuring warm and temperate. The average temperature is 16°C and average annual rainfall is 1676 mm, distributed throughout the year. Soil area is classified as Cambisol [15].

Soil samples were collected from a depth of 20 cm to the analysis (Table 1). Liming and fertilization were performed based on these.

### 2.2 Estimation of Diazotrophic Bacteria Community in Soil

It was estimated soil diazotrophic bacteria community [16]. A soil amount (10 g) was subjected to serial dilution. 0.5 ml of each dilution (from 10<sup>3</sup> to 10<sup>6</sup>) was inoculated in vials

**Table 1. Soil analysis from experimental area**

O.M g/dcm <sup>3</sup>	P mg/dcm <sup>3</sup>	pH CaCl <sub>2</sub>	H+Al	Al <sup>+3</sup>	K <sup>+</sup>	Ca <sup>+2</sup>	Mg <sup>+2</sup>	CTC	V
			Cmol <sub>c</sub> /dm <sup>3</sup>			%			
35.6	11.3	5.3	4,3	0.0	0.2	5.5	3,3	13.3	67.8

\*O.M. -Organic Matter

**Table 2. Treatments for analysis of *A. brasilense* inoculants efficiency to wheat development**

Treatments	Description
Co	Control - without inoculation and without Nitrogen topdressing fertilization.
CWI	Full Nitrogen topdressing fertilization without inoculation.
HCWI	Half of Nitrogen topdressing fertilization without inoculation.
CA5	Full Nitrogen topdressing fertilization with inoculant A containing Ab-V5.
HCA5	Half of Nitrogen topdressing fertilization with inoculant A containing Ab-V5.
CA56	Full Nitrogen topdressing fertilization with inoculant A containing Ab-V5 and Ab-V6
HCA56	Half of Nitrogen topdressing fertilization with inoculant A containing Ab-V5 and Ab-V6
CB56	Full Nitrogen topdressing fertilization with inoculant B containing Ab-V and Ab-V6.
HCB56	Half of Nitrogen topdressing fertilization with inoculant B containing Ab-V5 and Ab-V6.
WCA56	Inoculant A containing Ab-V5 and Ab-V6 without Nitrogen topdressing fertilization
WCB56	Inoculant B containing Ab-V5 and Ab-V6 without Nitrogen topdressing fertilization
WCA5	Inoculant A containing Ab-V5 without Nitrogen topdressing fertilization

containing 4.5 ml of the medium described in [16], with three replicates for each. It was checked the pellicle in the medium for diazotrophic bacteria presence.

### 2.3 Experimental Design and Parameters Analysis

The experimental design was in randomized block with twelve treatments (Table 2) and five replicates. The plot area was 24 m<sup>2</sup> (4 m x 6 m) with thirty-two rows spaced 0.17 m. The two inoculants tested (A and B) were at liquid formulation, containing 10<sup>8</sup> CFU (Colony Forming Units)/ml of *A. brasilense* strains, Ab-V5 and Ab-V6. The wheat cultivar was TBIO MESTRE, recommended for Santa Catarina state. Seeds were not treated with fungicides or insecticides. The seeds inoculation was performed according to the manufacturer's recommendation and manual planting. The top-dressing N fertilization was Urea (45%N) at 20 days after emergence (DAE), at 120 kg ha<sup>-1</sup> (full dosage) or 60 kg ha<sup>-1</sup> (half dosage). The experiment was carried out on July to November 2016.

Plants analyzed were taken from the plot center (1 m<sup>2</sup>). At 45 DAE, flag leaves N content were quantified [17]. At 115 DAE, it was performed the dry shoot weight, plant height, ear sizes, grains N contents and grain yield. The flag leaves and the grains were milled to perform the nitrogen content [17]. Plant height was measured from the base to the top, using graduated tape, as well as the ear size. Dry shoot weight was determined

after 72 h at 55°C. The productivity was determined by the grain weight at 13% humidity. The results were submitted to variance analysis (ANOVA) and when significant, compared by Scott-Knott's test, at 5% significance at Sisvar 5.6 software.

## 3. RESULTS AND DISCUSSION

### 3.1 Diazotrophic Bacteria Community in Soil

No pellicle was observed in the medium, even in the vials containing vials greater concentrations of cells, suggesting that the diazotrophic community was lower than 10<sup>4</sup> cells/g of soil. This result is consistent to those described by [1], in which the diazotrophic community in non-rhizospheric soil was around 10<sup>3</sup> cells/g of soil, before the development of sorghum roots. As the amount of *A. brasilense* cells in the inoculants was 10<sup>8</sup> CFU (Colony Forming Unit)/ml, applied directly on the seeds, that soil community did not influenced significantly on the results.

### 3.2 Wheat Development and Productivity Analysis

#### 3.2.1 Shoot dry weight, yield, ear size and plant height

There was no statistical difference for shoot dry weight, plant height, ear size and grain yield (Fig. 1). However, it is possible to observe that

for shoot dry weight (Fig. 1a), both inoculants, A and B, with the two strains, either N full or half dosage had more dry mass accumulation, highlighting CA56 that increased it by 17% compared to the control. The same could be observed for grain yield (Fig. 1b), which stood out HCA56. The association of inoculants, A and B, to the N top-dressing reached greater yields, ranging from 8% for CB56 to 17% for HCA56, respectively, when compared to the control. Among them, half N dosage reached more yields (12% for HCB56 and 17% for HCA56). The same effect was observed to ear size (Fig. 1c). For plant height, HCA56 and CA56 had higher individuals (Fig. 1d).


The significance absence to these parameters could be related to data variation. This results inconsistency was found in different studies [15,18,19,20]. Several reports did not observe a clear response to N fertilization and the inoculation with associative and/or endophytic diazotrophic bacteria. It suggested high edaphoclimatic influence and even, plant genotype variability. Spolaor, et al. [20] evaluated the association between topdressing N fertilization and two inoculants in IAC 125 popcorn cultivar and did not observe increase of productivity. Similar results were obtained in hybrid maize [21] and barley [19]. However, Mumbach, et al. [8] reported a positive relationship at inoculation and nitrogen fertilization for yield, number of ears/m<sup>2</sup> and

shoot dry matter to wheat. Araújo et al. [9] also reported increases in the number of green corn ears when bacteria were associated with gradual nitrogen dosage, suggesting other growth promotion mechanisms, such as auxin production, already described to rhizobacteria [12,13].


Despite the lack of statistical significance, the inoculants A and B, with the two strains, showed the highest means compared to treatments without inoculation or fertilization and that with 60 kg N/ha (half topdressing) (Fig. 1). This could be related to lower N amount available, especially as ammonium that, in high amounts, can reduce or inhibit nitrogenase activity [14]. Besides ammonium, glutamine, nitrate and nitrite can suppress the BFN and consequently many physiological processes, such as dry mass accumulation and yield. This indicates that the N amount in soil and organisms (plants or bacteria) regulates this process [14].

### 3.2.2 Nitrogen leaf and grain contents

There were statistical differences for leaves and grain N contents (Fig. 2). The leaf N amount was greater at the inoculant A, containing only strain Ab-V5, and half N dosage (HCA5), followed by the same inoculant A, with the two strains, and full top-dressing (CA56). The increases in N accumulation represented 109% and 87%, respectively, compared to the control (Fig. 2a).



**Fig. 1. Wheat growth and production parameters. (a) Shoot dry weight; (b) grain yield; (c) ear size; (d) Plant height**



**Fig. 2. Wheat N content in leaf (a) and in grain (b)**

Same letters do not differ at average test Scott Knott (5%)

The treatment HCA56 increased N accumulation in leaves, but less extent (28%). The other treatments did not differ from the control. Although, HCWI, CB56 and WCA5 accumulated less N in the leaves than the control.

The N grain content was significantly greater at inoculant B, containing the two strains, with no, half and full topdressing (WC56 - 51%, HCB56 - 76% and CB56 - 65%, respectively) and at inoculant A with the two strains and half topdressing (HCA56 - 62%) (Fig. 2b). All treatments had more N in the grains than the Co (Control).

The HCA5 and CA56 reached the highest N leaf levels, suggesting that the Ab-V5 strain was relevant to the N assimilation (Fig. 2a). However, despite the high N leaf content, the process of N translocation to the grains did not reflect the same efficiency (Fig. 2b). The N content in vegetative organs, such as leaves, is closely related to the translocation of sugars and nitrogen to grain [22]. Lana et al. [23] observed that the N leaf content was related to the increase in plant development parameters and N grain accumulation. However, they observed reduction at N leaf content for inoculation associated to nitrogen fertilization. These results were similar to those observed in this study. The N leaf accumulation was lower when complete nitrogen fertilization was associated to single Ab-V5 or to inoculant B, with the two strains (C5, HC5, CB56 and HB56 – Fig. 2a). The different responses to N accumulation, for grasses vegetative and production parameters seem to be related to genotypes [23] and even species [8,19,23]. Pereira et al. [24] observed that inoculation of Ab-V5 and Ab-V6 strains associated to 60 kg of N/ha increased grain yield and N content in grain. However, there was no

difference to the wheat shoot dry N content to the inoculated and uninoculated treatments.

For grain N accumulation, CWI, HCA56 CB56, HCB56 and WCB56 stood out, suggesting that the two strains in both inoculants were efficient to the translocation, as well as the 120 kg of N/ha fertilization. Vogel et al. [25] reviewing results of *A. brasilense* inoculation on wheat, connected positive results of productivity to the efficiency of biomass translocation to grains and the increase of photosynthetic activity. The authors also suggested detailed studies of those mechanisms. Current results reinforced the need to expand these investigations. Treatments that had the highest N content in grain (HCB56 and CB56) showed no leaf nitrogen accumulation significance, especially CB56 that had quantity below to the control. This suggested that higher N dosages, besides inhibiting the BNF can influence physiological processes because the microorganism can act like a drain, competing to the plant. Nevertheless, in general, associative diazotrophic bacteria, such as *A. brasilense*, can supply part of the N needed to the plant development. However, to maintain productivity indexes, it is necessary to associate BNF with N fertilization [8,19,20,23,24].

#### 4. CONCLUSION

Inoculation increased N content in wheat. The N leaf content was raised by the strain Ab-V5 of *A. brasilense* plus 60 Kg of N/ha (HCA5) and also by the strains Ab-V5 and Ab-V6 (inoculant A) plus 120 Kg of N/ha (CA56). The N grain content was raised by the strains Ab-V5 and Ab-V6 (inoculant B), with or without N fertilization (WCB56, HCB56 and CB56). These diazotrophic bacteria inoculation seems to be involved in N translocation, especially with lower fertilizing

supply, suggesting their potential as growth inducers, emphasizing the importance of further studies to confirm and understand the mechanisms involved.

## ACKNOWLEDGEMENT

To the agronomist Roberto de Almeida for technical and other supports. Mr. Silvio Ronaldo Nadal, for providing area on your farm “Fazenda do Tigre”. Grupo Vittia Biosoja Agrociências for financial support.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ. Análises de solo, planta e outros materiais (Boletim técnico 5). 2ª. ed. Porto Alegre: UFRGS - Departamento de Solos, Portuguese; 1995.
2. Notícias agrícolas. Bunge vê aumento da importação de trigo pelo Brasil em 2019 após quebra de safra; 2019. (Accessed 23 January 2019)  
Available: <https://www.noticiasagricolas.com.br/noticias/trigo/224825-bunge-ve-aumento-da-importacao-de-trigo-pelo-brasil-em-2019-apos-quebra-de-safra.html#.XEjhBFVKjIU>
3. Conab (Companhia Nacional de Abastecimento). Acompanhamento da safra brasileira de grãos. Observatório agrícola. Safra 2016/17 - Décimo levantamento, Brasília. 2017;4. (Accessed 26 October 2017)  
Available: [https://www.conab.gov.br/OlalaCMS/uploads/arquivos/17\\_07\\_12\\_11\\_17\\_01\\_boletim\\_graos\\_julho\\_2017.pdf](https://www.conab.gov.br/OlalaCMS/uploads/arquivos/17_07_12_11_17_01_boletim_graos_julho_2017.pdf)
4. SIDRA IBGE – LSPA (Levantamento Sistemático da Produção Agrícola). Tabela 1618- Área plantada, área colhida e produção, por ano da safra e produto; 2017. (Accessed 26 October 2017)  
Available: <https://sidra.ibge.gov.br/tabela/1618#resultado>
5. Wiethölter S. Fertilidade do solo e a cultura do trigo no Brasil. In: Pires JLM, et al. editors. Trigo no Brasil: Bases para produção competitiva e sustentável. Passo Fundo: Embrapa Trigo, Portuguese; 2011.
6. De Carvalho NLD, Zobot V. Nitrogênio: Nutriente ou poluente? Revista eletrônica em gestão, educação e tecnologia ambiental. Portuguese. 2012;6:960–974.  
Available: <http://dx.doi.org/10.5902/223611704671>
7. Quatrin MP, Olivo CJ, Simonetti GD, Bratz VF, De Godoy GL, Casagrande LGI. Response of dual-purpose wheat to nitrogen fertilization and seed inoculation with *Azospirillum brasilense*. Ciênc. Agrotec. 2019;43(3):1-10.  
Available: <http://dx.doi.org/10.1590/1413-7054201943027718>
8. Mumbach GL, Kotowski IE, Schneider FJA, Mallmann MS, Bonfada EB, Ortaça V, et al. Resposta da inoculação com *Azospirillum brasilense* nas culturas de trigo e de milho safrinha. Scientia Agraria. Portuguese. 2017;18(2):97-103.  
Available: <http://dx.doi.org/10.5380/rsa.v18i2.51475>
9. Araujo RM, Araujo ASF de, Nunes LAPL, Figueiredo, M do VB. Resposta do milho verde à inoculação com *Azospirillum brasilense* e níveis de nitrogênio. Ciência Rural. Portuguese. 2014;44(9):1556-1560.  
Available: <http://dx.doi.org/10.1590/0103-8478cr20130355>
10. Hungria M, Campo RJ, Mendes IC. A importância do processo de fixação biológica do nitrogênio para a cultura da soja: Componente essencial para a competitividade do produto brasileiro (Documentos 283). Londrina: Embrapa Soja; Portuguese; 2007.
11. Zilli JE, Campo RJ, Hungria M. Eficácia da inoculação de Bradyrhizobium em pré-semeadura da soja. Pesquisa Agropecuária Brasileira. Portuguese. 2010; 45(3):335-338.  
Available: <http://dx.doi.org/10.1590/S0100-204X2010000300015>
12. Olanrewaju OS, Glick BR, Babalola OO. Mechanisms of action of plant growth promoting bacteria. World Journal of Microbiology and Biotechnology. 2017; 33(197):1-16.  
Available: <http://dx.doi:10.1007/s11274-017-2364-9>
13. Bashan Y, De-Bashan L. How the plant growth-promoting bacterium *Azospirillum* promotes plant growth- A critical assessment. Advances in Agronomy. 2010; 108:77–136. (Accessed 25 December 2019)  
Available: <http://www.bashanfoundation.org/gmaweb/pdfs/mechanisms.pdf>

14. Carvalho TL, Balsemão-Pires E, Saraiva RM, Ferreira PC, Hemerly AS. Nitrogen signaling in plant interactions with associative and endophytic diazotrophic bacteria. *Journal of Experimental Botany*. 2014;65(19):5631–5642. Available: <https://doi.org/10.1093/jxb/eru319>
15. Hungria M, Campo RJ, Souza EM, Pedrosa FO. Inoculation with selected strains of *Azospirillum brasilense* and *A. lipoferum* improves yields of maize and wheat in Brazil. *Plant Soil*. 2010;331:413–425. Available: <http://dx.doi.org/10.1007/s11104-009-0262-0>
16. EMBRAPA. Centro Nacional de Pesquisa de Solos (Rio de Janeiro, RJ). Sistema brasileiro de classificação de solos. 2. ed. Rio de Janeiro: EMBRAPA-SPI, Portuguese; 2006. (Accessed 25 December 2019) Available: <https://www.agrolink.com.br/downloads/sistema-brasileiro-de-classificacao-dos-solos2006.pdf>
17. Estrada-De Los Santos P, Bustillos-Cristales R, Caballero-Mellado J. *Burkholderia*, a genus rich in plant-associated nitrogen fixers with wide environmental and geographic distribution. *Applied and Environmental Microbiology*. 2001;67(6):2790-2798. Available: <http://dx.doi.org/10.1128/AEM.67.6.2790-2798.2001>
18. Soares RS, Roesch LFW, Zanatta G, Camargo FA de O, Passaglia LMP. Occurrence and distribution of nitrogen fixing bacterial community associated with oat (*Avena sativa*) assessed by molecular and microbiological techniques. *Appl. Soil Ecol*. 2006;33:221-234. Available: <http://dx.doi.org/10.1016/j.apsoil.2006.01.001>
19. Barzotto GR, Lima SF, Santos OF, Piatí GL, Wassolowski CR. Adubação nitrogenada e inoculação com *Azospirillum brasilense* em cevada. *Nativa*. Portuguese. 2018;6(1):01-08. Available: <http://dx.doi.org/10.31413/nativa.v6i1.4611>
20. Spolaor LT, Gonçalves LSA, Dos Santos OJAP, de Oliveira ALM, Scapim CA, Bertagna FAB, Kuki MC. Bactérias promotoras de crescimento associadas à adubação nitrogenada de cobertura no desempenho agrônomo de milho pipoca. *Bragantia*. Portuguese. 2016;75(1):33-40. Available: <http://dx.doi.org/10.1590/1678-4499.330>
21. Dartora J, Guimarães VF, Marini D, Sander G. Adubação nitrogenada associada à inoculação com *Azospirillum brasilense* e *Herbaspirillum seropedicae* na cultura do milho. *Revista Brasileira de Engenharia Agrícola e Ambiental*. Portuguese. 2013;17(10):1023–1029. (Accessed 25 December 2019) Available: <http://www.scielo.br/pdf/rbeaa/v17n10/01.pdf>
22. Taiz L, Zeiger E. *Fisiologia Vegetal*. 3ª. ed. Porto Alegre: Artmed, Portuguese; 2004. Lana M, Dartora J, Marini D, Hann JE. Inoculation with *Azospirillum*, associated with nitrogen fertilization in maize. *Revista Ceres*. 2012;59(3):399-405. Available: <http://dx.doi.org/10.1590/S0034-737X2012000300016>
23. Pereira LC, Piana SC, Braccini AL, Garcia MM, Ferri GC, Felber PH et al. Rendimento do trigo (*Triticum aestivum*) em resposta a diferentes modos de inoculação com *Azospirillum brasilense*. *Revista de Ciências Agrárias*. Portuguese. 2017;240(1):105-113. Available: <https://doi.org/10.19084/RCA16089>
24. Vogel GF, Martinkoski L, Martins PJ, Bichel A. Desempenho agrônomo de *Azospirillum brasilense* na cultura do arroz: Uma revisão. *Revista em Agronegócios e Meio Ambiente*. Portuguese. 2013;6(3): 567-578.

© 2020 Balbinot et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/53471>